

The energy-carrying velocity and rolling of tachyons of unstable D-branes

Jin Hyun Chung^{a1} and W. S. L'Yi^{b2}

^a*Department of Liberal Arts, Chosun College of Science & Technology
290 Seosuk Dong, Dong-gu Kwang-ju City, Korea,*

^b*Department of Physics, Chungbuk National University,
Cheongju, Chungbuk 361-763, Korea.*

Abstract

It is shown that the tachyons which originate from unstable D-branes carry energy and momentum at the velocity $\beta = c^2/v$, where v is the phase velocity which is greater than c . For an observer who moves with the velocity β , tachyon is observed to be moving from one of the ground states of the tachyon potential to the potential hill. It is found that tachyon either passes over the hill or bounces back to the original ground state. Another possible solution is the case which is marginal to these, that is tachyon reaches to the top of the potential hill and stays there forever.

Nov.13, 2003

¹E-mail address: jhchung@mail.chosun-c.ac.kr

²E-mail address: wslyi@hep.chungbuk.ac.kr

1 Introduction

Tachyons, which abound in non-BPS D-branes of type IIA/IIB superstring theories or in bosonic strings, can be studied by using relatively simple effective action[1]. The tachyon potential $V(T)$, which is an even function of the tachyon field T , has a run-away form: it is maximum at $T = 0$, and minimum at $T \rightarrow \pm\infty$. This means that if the initial string state is the tachyonic state around the unstable false vacuum $T \approx 0$, all the energy, because of the fact that the full string states contain various field modes, will eventually be distributed among various bulk modes.

This process of rolling of tachyons and tachyon condensations are discussed by various authors[2]. Recently it is pointed out that fluctuations around a static kink solution of D_p -brane produces the Dirac-Born-Infeld action of a BPS D_{p-1} -brane[3]. It is also found[4] that the only static finite energy solution of tachyon field which depends only on one of the spatial coordinate x is $T(x) = Ax$, where $A \rightarrow \infty$.

In this paper, we investigate the traveling wave solutions of the tachyon field which propagate along x . It is found that the phase velocity v of tachyon wave may have any value except the speed of light c . As usual, the phase velocity is quite different from the mechanical velocity¹ at which the energy-momentum is carried. If $v < c$, they are the same when one assumes that the wave amplitude A becomes very large. On the other hand, if $v > c$, the mechanical velocity is $\beta = c^2/v$. Furthermore, the invariant mass $m = \sqrt{E^2 - p^2 c^2}/c^2$ is always real. This important fact is pointed out in[5] where boundary state idea is used under different perspective. It means that tachyonic states of unstable D-branes can be interpreted as usual matter states moving at the mechanical velocity β . This energy distribution in space may be related to the final bulk states of

¹The usual terminology for the velocity at which waves carry the energy-momentum is the group velocity. But in this tachyonic case whose highly non-linear equation of motion does not admit the simple rule of linear superposition, it is more adequate to use the term mechanical velocity instead of group velocity.

strings.

For an observer who moves with the velocity β , tachyon is observed to be rolling from one of the ground states of the tachyon potential to the potential hill. It is found that there are three different types of motion. The first is that tachyon passes over the hill and rolls to another ground state. The second one is the case that it can not cross over the hill and bounces back to the original ground state. The last possibility is marginal to these, and tachyon reaches to the top of the potential hill and stays there forever.

In section 2, a propagating wave solution for the tachyon is obtained. It has either oscillating form or kink form. The oscillating solution is studied in section 3, and the kink solution is investigated in section 4. Physical interpretation of the solution is given in section 5. The conclusion is given in section 6.

2 Propagating wave solution for the tachyon field

Consider the following effective action of the tachyon field T on a non-BPS D_p -brane,

$$S = - \int d^{p+1}x V(T) \sqrt{-\det(\eta_{\mu\nu} + \partial_\mu T \partial_\nu T)}. \quad (1)$$

Here, $\eta_{\mu\nu} = \text{diag}(-1, 1, \dots, 1)$, and $\hbar = c = \alpha' = 1$. For the potential of the tachyon field, we assume the following form[6, 7]

$$V(T) = \frac{\tau_p}{\cosh(\frac{T}{T_0})}, \quad (2)$$

where τ_p is the brane tension, and T_0 is a constant which is $\sqrt{2}$ for non-BPS D-branes in superstrings, and 2 for bosonic strings.

To solve the Lagrangian equation of motion, we introduce a new variable $\mathcal{T}$ defined by[7, 8]

$$\mathcal{T} = T_0 \sinh \frac{T}{T_0}. \quad (3)$$

The Lagrangian in terms of $\mathcal{T}$, which is given by

$$\mathcal{L} = -\frac{\tau_p}{1 + \frac{\mathcal{T}^2}{T_0^2}} \sqrt{1 + \frac{\mathcal{T}^2}{T_0^2} + \partial_\mu \mathcal{T} \partial^\mu \mathcal{T}}, \quad (4)$$

does not contain any transcendental function. The corresponding equation of motion is given by the following polynomial form of $\mathcal{T}$, $\partial_\mu \mathcal{T}$ and $\partial_\mu \partial_\nu \mathcal{T}$:

$$(1 + \frac{\mathcal{T}^2}{T_0^2})(\partial_\mu \partial^\mu \mathcal{T} + \frac{\mathcal{T}}{T_0^2}) + \partial_\mu \mathcal{T} \partial^\mu \mathcal{T} \partial_\nu \partial^\nu \mathcal{T} - \partial_\mu \mathcal{T} \partial_\nu \mathcal{T} \partial^\mu \partial^\nu \mathcal{T} = 0. \quad (5)$$

We observe the following facts that in each term of this equation, when expanded, ∂_μ appears even times while $\mathcal{T}$ odd times. This means that for a given solution, one may obtain another solution by changing either the sign of a particular x^μ or/and $\mathcal{T}$.

It is known[3] that for the extreme *static* kink solution such as $T \rightarrow \pm\infty$ depending on the sign of x^p , the resulting configuration is a stable D_{p-1} -brane at $x = 0$. In this paper we assume that $\mathcal{T}$, or equivalently T , depends on both $x = x^p$ and t . In this case, (5) can be written as

$$(1 + \frac{\mathcal{T}^2}{T_0^2})(\partial_x^2 - \partial_t^2 + \frac{1}{T_0^2})\mathcal{T} + 2\partial_x \mathcal{T} \partial_t \mathcal{T} \partial_x \partial_t \mathcal{T} - (\partial_x \mathcal{T})^2 \partial_t^2 \mathcal{T} - (\partial_t \mathcal{T})^2 \partial_x^2 \mathcal{T} = 0. \quad (6)$$

It is easy to show that it has the following traveling wave solution,

$$\mathcal{T} = \frac{T_0}{2} \left\{ A e^{ik(x-vt)} + B e^{-ik(x-vt)} \right\}, \quad (7)$$

where A , B and v are all integration constants, and k is given by

$$k = \frac{1}{\sqrt{1-v^2}} \frac{1}{T_0}. \quad (8)$$

If $|v| < 1$, k is real and the solution is oscillatory. This is discussed in section 3. On the other hand, if $|v| > 1$, one may replace ik by κ to obtain a kink solution. It is discussed in section 4.

3 Traveling sinusoidal solution

Consider the case $|v| < 1$. For the convenience of computation we rewrite (7) in the following form

$$\mathcal{T} = T_0 A \cos k(x - vt). \quad (9)$$

To understand the physical meaning of the phase velocity v , we calculate the energy and momentum densities of the field configuration. The relevant energy-momentum tensor $T^\mu{}_\nu$ is

$$T^\mu{}_\nu = -\frac{\partial \mathcal{L}}{\partial \partial_\mu T} \partial_\nu T + \delta^\mu{}_\nu \mathcal{L}, \quad (10)$$

where the Lagrangian expressed in terms of $T(x, t)$ is given by

$$\mathcal{L} = -V(T) \sqrt{1 + (\partial_x T)^2 - (\partial_0 T)^2}. \quad (11)$$

From this one can read off the following energy density $\rho = T^{00}$ and momentum density $P = T^{p0}$,

$$\rho = \frac{V(T)}{\sqrt{1 + (\partial_x T)^2 - (\partial_0 T)^2}} \{1 + (\partial_x T)^2\}, \quad (12)$$

$$P = \frac{V(T)}{\sqrt{1 + (\partial_x T)^2 - (\partial_0 T)^2}} \partial_x T \partial_0 T. \quad (13)$$

After non-trivial but straight forward calculations it can be shown that

$$\frac{V(T)}{\sqrt{1 + (\partial_x T)^2 - (\partial_0 T)^2}} = \frac{\tau_p}{\sqrt{1 + A^2}}. \quad (14)$$

On the other hand, by using the defining equation (3) and the solution (9), one may prove that

$$(\partial_x T)^2 = \frac{(\partial_x \mathcal{T})^2}{1 + \frac{\mathcal{T}^2}{T_0^2}} \quad (15)$$

$$= k^2 T_0^2 \frac{A^2 \sin^2 k(x - vt)}{1 + A^2 \cos^2 k(x - vt)}. \quad (16)$$

That is, ρ and P , for this traveling sinusoidal solution, are

$$\rho = \frac{\tau_p}{\sqrt{1+A^2}} \left\{ 1 + \frac{1}{1-v^2} \frac{A^2 \sin^2 k(x-vt)}{1+A^2 \cos^2 k(x-vt)} \right\}, \quad (17)$$

$$P = \frac{\tau_p}{\sqrt{1+A^2}} \frac{v}{1-v^2} \frac{A^2 \sin^2 k(x-vt)}{1+A^2 \cos^2 k(x-vt)}. \quad (18)$$

When $A = 0$, one has the trivial unstable solution, $T = 0$. In this case, $\rho = \tau_p$ and $P = 0$ as expected.

For general A , the energy density is oscillatory at the string length scale T_0 . The average behaviour of it can be obtained by using the relation,

$$\frac{1}{2\pi} \int_0^{2\pi} \frac{a \sin^2 \theta}{1+a \cos^2 \theta} d\theta = \sqrt{1+a} - 1. \quad (19)$$

That is, the spatial average of ρ is

$$\bar{\rho} = \frac{\tau_p}{1-v^2} \left(1 - \frac{v^2}{\sqrt{1+A^2}} \right). \quad (20)$$

For large A , it becomes

$$\bar{\rho} \simeq \frac{\tau_p}{1-v^2}. \quad (21)$$

It can be interpreted in the following way. The energy stored in length L is $E_L = \bar{\rho}L$. Here, one should be careful that it is not L but $L_0 = \gamma L$ which is Lorentz invariant, where $\gamma = 1/\sqrt{1-v^2}$. It means that it is more appropriate to express E_L in terms of L_0 ,

$$E_L = \gamma \tau_p L_0. \quad (22)$$

This shows that the (rest) mass density is $\rho_m = \tau_p$.

Similarly, the average momentum density is

$$\bar{P} = \frac{\tau_p v}{1-v^2} \left(1 - \frac{1}{\sqrt{1+A^2}} \right). \quad (23)$$

This means that, for large A , the momentum in length L is

$$P_L = \bar{P}L \quad (24)$$

$$= \gamma \tau_p v L_0. \quad (25)$$

Combining this with (22) one gets $P_L = vE_L$. It shows that v is in fact the velocity at which the energy is propagated. Similarly we have $E_L^2 - P_L^2 = m_L^2$, where $m_L = \tau_p L_0$ is the mass stored in the invariant length L_0 .

4 Traveling kink solution

Consider the case $|v| > 1$. In this case, by replacing ik of (7) with κ we obtain the following solution,

$$\mathcal{T} = \frac{T_0}{2} \left\{ A e^{\kappa(x-vt)} + B e^{-\kappa(x-vt)} \right\}. \quad (26)$$

Here,

$$\kappa = \frac{1}{\sqrt{v^2 - 1}} \frac{1}{T_0}. \quad (27)$$

The corresponding energy and momentum densities which can be determined from (12) and (13) are

$$\rho = \frac{\tau_p}{\sqrt{1+AB}} \left\{ 1 + (\partial_x T)^2 \right\}, \quad (28)$$

$$P = \frac{\tau_p}{\sqrt{1+AB}} (\partial_x T)^2 v. \quad (29)$$

By using the same technique employed in section 3, one may prove that

$$(\partial_x T)^2 = \kappa^2 T_0^2 \frac{A^2 e^{2\kappa(x-vt)} + B^2 e^{-2\kappa(x-vt)} - 2AB}{A^2 e^{2\kappa(x-vt)} + B^2 e^{-2\kappa(x-vt)} + 2AB + 4}. \quad (30)$$

That is,

$$\rho = \frac{\tau_p}{\sqrt{1+AB}} \left\{ 1 + \kappa^2 T_0^2 \frac{A^2 e^{2\kappa(x-vt)} + B^2 e^{-2\kappa(x-vt)} - 2AB}{A^2 e^{2\kappa(x-vt)} + B^2 e^{-2\kappa(x-vt)} + 2AB + 4} \right\}, \quad (31)$$

$$P = \frac{\tau_p}{\sqrt{1+AB}} \kappa^2 T_0^2 \frac{A^2 e^{2\kappa(x-vt)} + B^2 e^{-2\kappa(x-vt)} - 2AB}{A^2 e^{2\kappa(x-vt)} + B^2 e^{-2\kappa(x-vt)} + 2AB + 4} v. \quad (32)$$

To have real ρ and P , it is required that $AB > -1$. We consider $AB > 0$, and $0 > AB > -1$, $AB = 0$ cases separately.

Case 1) $AB > 0$.

In this case, one may assume that $A = B$. It is always possible by properly choosing the origin of x . The solution has the following form

$$\mathcal{T} = T_0 A \cosh \kappa(x - vt). \quad (33)$$

The energy density ρ which can be read off from (31) is

$$\rho = \frac{\tau_p}{\sqrt{1 + A^2}} \left\{ 1 + \frac{1}{v^2 - 1} \frac{A^2 \sinh^2 \kappa(x - vt)}{A^2 \cosh^2 \kappa(x - vt) + 1} \right\}. \quad (34)$$

This ρ is almost constant except at the region of a small dimple of size T_0 .

By using the relation

$$\int \frac{a \sinh^2 x}{1 + a \cosh^2 x} dx = x - \sqrt{1 + a} \tanh^{-1} \left\{ \frac{\tanh x}{\sqrt{1 + a}} \right\} + \text{const.}, \quad (35)$$

one can show that the stored energy in the region $-\frac{L}{2} < x < \frac{L}{2}$ of length L is the following,

$$E_L = \frac{v^2}{v^2 - 1} \frac{\tau_p}{\sqrt{1 + A^2}} L - \frac{2\tau_p}{v^2 - 1} \tanh^{-1} \left(\frac{\tanh \frac{L}{2}}{\sqrt{1 + A^2}} \right). \quad (36)$$

That is, the energy density $\bar{\rho} = \lim_{L \rightarrow \infty} E_L/L$ effectively becomes

$$\bar{\rho} = \frac{\tau_p}{\sqrt{1 + A^2}} \frac{1}{1 - \beta^2}, \quad (37)$$

where $\beta = 1/v$.

Similarly, we have

$$\bar{P} = \frac{\tau_p}{\sqrt{1 + A^2}} \frac{\beta}{1 - \beta^2}. \quad (38)$$

It is interesting that the momentum density $\bar{P}$ vanishes if $\beta \rightarrow 0$, or equivalently if $v \rightarrow \infty$. This gives us a hint that v is not the mechanical velocity. In fact the relation $\bar{P} = \beta \bar{\rho}$ shows that it is not v but β which carries energy. This means that tachyon can be interpreted as a usual matter which moves at the mechanical velocity β . Similar analysis done in section 3 may be employed here to show that the mass density is

$$\rho_m = \frac{\tau_p}{\sqrt{1 + A^2}}. \quad (39)$$

Case 2) $0 > AB > -1$.

By a suitable choice of the origin of x one may assume that $B = -A$, where $0 < A^2 < 1$. Then, the solution has the following form,

$$\mathcal{T} = T_0 A \sinh \kappa(x - vt). \quad (40)$$

The corresponding energy density ρ is

$$\rho = \frac{\tau_p}{\sqrt{1 - A^2}} \left\{ 1 + \frac{1}{v^2 - 1} \frac{A^2 \cosh^2 \kappa(x - vt)}{A^2 \sinh^2 \kappa(x - vt) + 1} \right\} \quad (41)$$

As the case 1), ρ has a small dimple of size T_0 . By using the relation

$$\int \frac{a \cosh^2 x}{1 + a \sinh^2 x} dx = x - \sqrt{1 - a} \tanh^{-1} \left\{ \sqrt{1 - a} \tanh x \right\} + \text{const.}, \quad (42)$$

it can be shown that the total energy and momentum stored in the region $-\frac{L}{2} < x < \frac{L}{2}$ of length L are

$$E_L = \frac{1}{1 - \beta^2} \frac{\tau_p}{\sqrt{1 - A^2}} L - \frac{2\tau_p \beta^2}{1 - \beta^2} \tanh^{-1} \left(\sqrt{1 - A^2} \tanh \frac{L}{2} \right), \quad (43)$$

$$P_L = \frac{\beta}{1 - \beta^2} \frac{\tau_p}{\sqrt{1 - A^2}} L - \frac{2\tau_p \beta}{1 - \beta^2} \tanh^{-1} \left(\sqrt{1 - A^2} \tanh \frac{L}{2} \right). \quad (44)$$

That is, the energy and momentum densities, when $L \rightarrow \infty$, effectively become

$$\bar{\rho} = \frac{1}{1 - \beta^2} \frac{\tau_p}{\sqrt{1 - A^2}}, \quad (45)$$

$$\bar{P} = \frac{\beta}{1 - \beta^2} \frac{\tau_p}{\sqrt{1 - A^2}}. \quad (46)$$

These show that the tachyon can in fact be interpreted as a usual matter of mass density

$$\rho_m = \frac{\tau_p}{\sqrt{1 - A^2}} \quad (47)$$

moving at the velocity $\beta = 1/v$.

Case 3) $AB = 0$.

We assume that $B = 0$. The other case, that is $A = 0$, can be obtained from the results of this case by replacing $\kappa \rightarrow -\kappa$. The corresponding solution is given by

$$\mathcal{T} = Ae^{\kappa(x-vt)}. \quad (48)$$

The energy and momentum densities are

$$\rho = \tau_p \left\{ 1 + \frac{1}{v^2 - 1} \frac{A^2 e^{2\kappa(x-vt)}}{A^2 e^{2\kappa(x-vt)} + 4} \right\}, \quad (49)$$

$$P = \tau_p \left\{ \frac{1}{v^2 - 1} \frac{A^2 e^{2\kappa(x-vt)}}{A^2 e^{2\kappa(x-vt)} + 4} \right\} v. \quad (50)$$

$$(51)$$

These mean that the energy and momentum densities under the $A \rightarrow \infty$ limits become

$$\rho = \frac{\tau_p}{1 - \beta^2}, \quad (52)$$

$$P = \frac{\tau_p}{1 - \beta^2} \beta. \quad (53)$$

This means that it also describes the usual matter of the mass density $\rho_m = \tau_p$ moving at the velocity $\beta = 1/v$.

5 Interpretation of the solutions

Consider the traveling sinusoidal solution (9) first. For an observer with coordinates (x', t') who moves at the velocity v with respect to the frame (x, t) , the solution can be written as

$$\mathcal{T} = T_0 A \cos \frac{x'}{T_0}. \quad (54)$$

This represents an infinite lattice of brain-antibrane system[9].

For the traveling kink solution, consider (26). For an observer with coordinates (x', t') who moves with velocity β , phase $\kappa(x - vt)$ can be written as $-\frac{t'}{T_0}$, where we used

$$\frac{x - vt}{\sqrt{v^2 - 1}} = -t'. \quad (55)$$

This means that for kink solution it can be written as

$$\mathcal{T} = \begin{cases} T_0 A \cosh(-\frac{t'}{T_0}) & \text{for Case 1,} \\ T_0 A \sinh(-\frac{t'}{T_0}) & \text{for Case 2,} \\ T_0 A e^{-\frac{t'}{T_0}} & \text{for Case 3.} \end{cases} \quad (56)$$

It is clear that Case 2 represents motion of tachyon which moves from one ground state of the the potential at $t' \rightarrow -\infty$ to the other ground state at $t' \rightarrow \infty$, thus crossing over the potential hill. For the Case 1, tachyon does not cross over the potential hill, and begins to bounce back to the original vacuum at $t' = 0$. The Case 3 is marginal to these, and tachyon approaches to the maximum of the potential as $t' \rightarrow \infty$.

6 Conclusion

The traveling wave solution of the Sen's tachyon action, either in the oscillating form or kink form, represents usual matter that moves at the velocity β which is less than c . If the phase velocity v is less than c , the tachyon field oscillates in space. In this case, $\beta = v$ and the mass density is τ_p .

If, on the other hand, the phase velocity of the tachyon field v is greater than the speed of light c , tachyon field forms a kink. In this case, the velocity at which energy is carried is $\beta = c^2/v$. This corresponds to the rolling of tachyons. The corresponding mass density is $\tau_p/\sqrt{1+A^2}$, $\tau_p/\sqrt{1-A^2}$ or τ_p depending on the precise form of the solution. This energy-momentum may be distributed among various modes of strings.

Acknowledgement

The author would like to thank Paul Frampton for his interests and critical comments on this paper, and for his hospitality during the author's stay in UNC, Chapel Hill.

References

- [1] A. Sen, JHEP **9910**, 008 (1999) [arXiv:hep/9909062]; Mod. Phys. Lett. A**17**, 1797 (2002), [arXiv:hep-th/0204143]; arXiv:hep-th/0209122; M. R. Garousi, Nucl. Phys. B **584**, 284 (2000) [arXiv:hep-th/0003122]; Nucl. Phys. B **647**, 117 (2002) [arXiv:hep-th/0209068]; E. A. Bergshoeff, M. de Roo, T. C. de Wit, E. Eyras and S. Pand, JHEP **0005**, 009 (2000) [arXiv:hep-th/0003221]; J. Kluson, Phys. Rev. D **62**, 126003 (2000) [arXiv:hep-th/0004106].
- [2] A. Sen, JHEP **0204**, 048 (2002) [arXiv:hep-th/0203211]; JHEP **0207**, 065 (2002) [arXiv:hep-th/0203265]; Mod. Phys. Lett. A **17**, 1797 (2002) [arXiv:hep-th/0204143]; K. Larsen, A. Naqvi and S. Terashima, arXiv:hep-th/0212248; James M. Cline and Hassan Firouzjahi, arXiv:hep-th/0301101; S. Sugimoto and S. Terashima, JHEP **0207**, 025 (2002) [arXiv:hep-th/0205085]; J. A. Minahan, JHEP **0207**, 030 (2002) [arXiv:hep-th/0205098]; G. Gibbons, K. Hashimoto and P. Yi, JHEP **0209**, 061 (2002) [arXiv:hep-th/0209034]; D. Kutasov, M. Marino and G. W. Moore, JHEP **0010**, 045 (2000) [arXiv:hep-th/0009148]; C. Vafa, arXiv:hep-th/0111051; J. R. David, M. Gutperle, M. Headrick and S. Minwalla, JHEP **0202**, 041 (2002) [arXiv:hep-th/0111212].
- [3] A. Sen, arXiv:hep-th/0303057
- [4] Ph. Brax, J. Mourad and D. A. Steer, arXiv:hep-th/0304197.
- [5] Soo-Jong Rey and Shigeki Sugimoto, Phys. Rev. D **67**, 086008 (2003) [arXiv:hep-th/0301049]; Soo-Jong Rey and Shigeki Sugimoto, arXiv:hep-th/0303133.
- [6] C. Kim, H. B. Kim, Y. Kim, and O-K. Kwon, JHEP **0301**, 008 (2003) [arXiv:hep-th/0301076]; F. Leblond and A. W. Peet, arXiv:hep-th/0303035; N. Lambert, H. Liu and J. Maldacena, arXiv:hep-th/0303139.

- [7] David Kutasov and Vasilis Niarchos, arXiv:hep-th/0304045.
- [8] Kazumi Okuyama, arXiv:hep-th/0304108.
- [9] C. Kim, Y. Kim, O. K. Kwon, and C. O. Lee, arXiv:hep-th/0305092, Chanju Kim, Yoonbai Kim, O-Kab Kwon, and Piljin Yi, arXiv:hep-th/0307184